Protective Effects of Ursolic Acid on Osteoblastic Differentiation via Activation of IER3/Nrf2

Sang-im Lee[†]

Department of Dental Hygiene, College of Health Sciences, Dankook University, Cheonan 31116, Korea

Background: Oxidative stress is known to be associated with the pathogenesis of many inflammatory diseases, including periodontitis. Ursolic acid is a pentacyclic triterpenoid with antimicrobial, antioxidative, and anticancer properties. However, the role of ursolic acid in the regulating of osteogenesis remains undetermined. This study was aimed to elucidate the crucial osteogenic effects of ursolic acid and its ability to inhibit oxidative stress by targeting the immediate early response 3 (IER3)/nuclear factor erythroid 2-related factor 2 (Nrf2) pathway.

Methods: Cell proliferation was determined using water-soluble tetrazolium salt assay, cell differentiation was evaluated by alkaline phosphatase (ALP) activity, and formation of calcium nodules was detected using alizarin red S stain. Generation of reactive oxygen species (ROS) was determined using DCFH-DA fluorescence dye in hydrogen peroxide (H₂O₂)-treated MG-63 cells. Expression levels of IER3, Nrf2, and heme oxygenase-1 (HO-1) were analyzed using western blot analysis.

Results: Our results showed that ursolic acid up-regulated the proliferation of osteoblasts without any cytotoxic effects, and promoted ALP activity and mineralization. H₂O₂-induced ROS generation was found to be significantly inhibited on treatment with ursolic acid. Furthermore, in H₂O₂-treated cells, the expression of the early response genes: IER3, Nrf2, and Nrf2-related phase II enzyme (HO-1) was enhanced in the presence of ursolic acid.

Conclusion: The key findings of the present study elucidate the protective effects of ursolic acid against oxidative stress conditions in osteoblasts via the IER3/Nrf2 pathway. Thus, ursolic acid may be developed as a preventative and therapeutic agent for mineral homeostasis and inflammatory diseases caused due to oxidative injury.

Key Words: Immediate early response 3 protein, NF-E2-related factor 2, Osteoblast, Oxidative stress, Periodontal disease

Introduction

Oxidative stress is well-known for disturbing the equilibrium between the production of reactive oxygen species (ROS) and the ability of the biological system to promote antioxidant defenses¹⁾. ROS affect all biomolecules such as carbohydrates, membrane lipids, proteins, and nucleic acids, and thus, caused detrimental events such as physiological adaptation phenomena and disruption in the regulation of intracellular signal transduction²⁾. Consequently, oxidative stress has been associated with the pathology of many inflammatory diseases³⁾, including cancer, atherosclerosis, hypertension, rheumatoid arthritis⁴⁾, and osteoporosis⁵⁾. In addition, induction of oxidative stress is a

potential mechanism that provokes periodontal disease which affects systemic health⁶⁾.

Periodontal disease, a chronic inflammatory disease which is induced initiated by bacterial infection, degrades the supporting tissues of the teeth. In its advanced stage, periodontitis causes a severe immune reaction in the periodontal tissues leading to tooth loss in adults and potential systemic effects^{7,8)}. Recently, many studies have reported that ROS contributes in inducing oxidative stress and the pathogenesis of periodontal disease in humans^{9,10)}. Importantly, studies have shown that oxidative stress decreases the osteogenic potential of mesenchymal stem cells, which are the progenitors of osteoblasts, adipocytes, and chondrocytes¹¹⁾. Furthermore, it has been reported to

Received: August 6, 2019, Revised: August 28, 2019, Accepted: September 3, 2019

eISSN 2233-7679

[†]Correspondence to: Sang-im Lee, <https://orcid.org/0000-0003-2635-6876>

Department of Dental Hygiene, College of Health Sciences, Dankook University, 119 Dandae-ro, Dongnam-gu, Cheonan 31116, Korea
Tel: +82-41-550-1492, E-mail: hanjumuck@dankook.ac.kr

Copyright © The Korean Society of Dental Hygiene Science.

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

cause periodontitis by upregulating osteoclastic bone resorption, which degrades the connective tissue and alveolar bone supporting the teeth during inflammatory response¹²⁾. Several evidences have suggested that oxidative stress, as a critical mediator of pathophysiological response, is a negative regulator for bone hemostasis. It has been well documented that ROS regulates various signaling pathways that are involved in bone remodeling such as Wnt/ β -catenin signaling¹³⁾, mitogen-activated protein kinase signaling¹⁴⁾ and the nuclear factor erythroid 2-related factor 2 (Nrf2) signaling pathways¹⁵⁾.

Ursolic acid (3 β -hydroxy-urs-12-en-28-oic acid) is known as a natural pentacyclic triterpenoid, which has been used traditionally as a herbal medicine in Eastern Asia for treating many clinical diseases¹⁶⁾. Ursolic acid is found in various fruits, medicinal herbs, and other plants. The natural compounds has been reported to possess several biological effects, including antioxidant, anti-inflammatory, and anticancer activity¹⁷⁾. Despite various beneficial effects reported in the clinical field, the underlying mechanisms of its action remain unknown.

Consequently, this study was performed to investigate the beneficial effects of ursolic acid and its underlying mechanisms of the action of ursolic acid against hydrogen peroxide (H₂O₂)-induced oxidative stress in the pathogenic conditions of periodontal disease.

Materials and Methods

1. Cell culture

MG-63 osteoblasts were obtained from the American Type Culture Collection (ATCC, Manassas, VA, USA) and cultured as per the recommended instructions. Dulbecco's Modified Eagle Medium (DMEM), fetal bovine serum (FBS), trypsin/ethylenediaminetetraacetic acid, and penicillin/streptomycin were from Gibco (Life Technologies, Grand Island, NY, USA). Cells were cultured in DMEM supplemented with 10% FBS, and penicillin/streptomycin (100 μ g/ml) in a humidified atmosphere containing 5% CO₂ at 37°C. Further, cells were plated in 24-well culture plates at a density of 1 $\times$ 10⁴ cells/well. For osteoblast differentiation, cells were treated with osteogenic medium (OS) containing 10% FBS, 50

μ g/ml ascorbic acid, and 10 mM β -glycerophosphate. OS was replaced every 3 days. To induced oxidative stress, cells were supplemented with fresh medium and treated with different concentrations of H₂O₂ (100 μ M) for 24 hours or 10 days.

2. Cell proliferation assays

Toxicity of ursolic acid (Sigma-Aldrich Corp., St. Louis, MO, USA) on MG-63 cells was assessed by water-soluble tetrazolium salt (WST-1) cell proliferation assays (Sigma-Aldrich Corp.). Briefly, after plating the cells in a 96-well plate (5 $\times$ 10³ cells/well) for 24 hours, they were treated with different concentrations of H₂O₂ (100 μ M). Then, WST-1 was added each day for 7 days (10 μ L/per well) and 4 hours in the dark humid conditions. Microplates were gently mixed by shaking for 1 minute and absorbance was measured at 450 nm using the Bio-Rad 680 plate-reader (Bio-Rad Labs, Hercules, CA, USA).

3. Reactive oxygen species measurement

ROS levels were detected using 2', 7'-dichlorofluorescein diacetate (DCFH-DA; Beyotime, Shanghai, China) according to the manufacturer's instructions. Cells were treated with 100 μ M hydrogen peroxide in the presence or absence of ursolic acid, and incubated for 1 hour. After washing with phosphate buffer saline (PBS), the cells were stained with 10 μ M DCFH-DA in the dark. Further, the cells were harvested and measured by flow cytometry (BD Biosciences, San Jose, CA, USA) with an excitation at 490 nm and an emission at 530 nm. The generation of intracellular ROS was then quantified.

4. Alkaline phosphatase activity assay

For determining alkaline phosphatase activity (ALP) activity, the cells were plated in 96-well plates at 1 $\times$ 10⁴ cells/well and cultured in OS for 14 days. ALP activity was analyzed by following the protocol recommended in ALP fluorometric assay kit (BioVision, Milpitas, CA, USA). Absorbance was measured at 405 nm on a spectrophotometer.

5. Alizarin red S stain

Cell culture plates were washed twice with PBS and fixed at room temperature with 4% formaldehyde for 10 min. The plates were then washed gently three times with dH₂O, stained at room temperature with 1% Alizarin Red S (Sigma-Aldrich) for 10 minutes, and visualized using a fluorescence microscope (Olympus IX73; Olympus Corporation, Tokyo, Japan).

6. Western blot analysis

PRO-PREPTM (iNtRON Biotechnology, Seongnam, Korea) was used to lyse cells, and bicinchoninic acid assay (Pierce, Appleton, WI, USA) was performed to determine the protein expression levels. Total protein (30 µg per lane) was run on a 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis under reducing conditions, and then proteins were transferred onto a polyvinylidene difluoride membrane. The membranes were blocked with 5% non-fat milk in PBS containing 0.1% Tween-20 (PBS-T) for 1 hour, rinsed with PBS-T, and incubated with primary antibodies against (1:1,000 dilution) overnight at 4°C. Antibody for β -actin was purchased from Sigma-Aldrich, whereas antibodies for immediate early response 3 (IER3), Nrf2, and heme oxygenase-1 (HO-1) were purchased from Cell Signaling (Lake Danvers, MA, USA). Secondary antibodies (horseradish peroxidase-conjugated) were incubated for 30 minutes, blots were visualized by chemiluminescence,

and exposed to X-ray film. Image-ProPlus 5.0 software (Media Cybernetics Inc., Rockville, MD, USA) was used for quantification.

7. Statistical analysis

Data are presented as mean $\pm$ standard deviation from a minimum of three replicates. The results were statistically analyzed by paired Student's t-test and ANOVA using IBM SPSS ver. 25.0 software (IBM Corp., Armonk, NY, USA).

Results

1. Ursolic acid induced cell proliferation and differentiation in osteoblasts

At first, we determined the effect of ursolic acid on cell proliferation in MG-63 cells by using the WST-1 assays. The cells were treated with ursolic acid (0.5 µM, 1 µM, 5 µM, 10 µM) for 24 hours. The results showed that ursolic acid promoted higher proliferation rates compared to the untreated controls without a cytotoxic effect (Fig. 1A). ALP has been shown to be secreted by differentiated osteoblasts, which is correlated to the early stages of osteogenic differentiation¹¹⁾. Our data showed that ursolic acid promotes the ALP activity in a dose-dependent as well as time-dependent manner for 10 days. Remarkably, we observed that 5 µM ursolic acid treatment group showed a 242.4 $\pm$ 12.1% increase in ALP activity compared to the

Fig. 1. Effects of ursolic acid on cell proliferation and differentiation of osteoblasts. (A) Cells were treated with various concentration of ursolic acid for 24 hours. Cell proliferation was evaluated by water-soluble tetrazolium salt assay. Osteoblastic differentiation was assessed by alkaline phosphatase (ALP) activity (B) and alizarin red S stain (C). Values are mean $\pm$ standard deviation of three experiments. * p < 0.05 vs. control group.

activity shown by the untreated control group ($p < 0.05$) (Fig. 1B). Furthermore, formation of mineralized nodules is one of the markers for the osteoblastic maturation. Alizarin red S staining showed that mineralized nodule formation after the osteoblasts were cultured in the OS with ursolic acid for 10 days. We found that ursolic acid up-regulates also dose-dependently increase extracellular matrix mineralization in a dose-dependent manner (Fig. 1C). Thus, our results suggested that ursolic acid promotes osteogenic differentiation.

2. Ursolic acid inhibited H_2O_2 -induced oxidative stress in osteoblasts

To further examine whether ursolic acid contributes in regulating oxidative stress and the differentiation in MG-63 cells during osteogenic-induction, the cells were treated with $100 \mu M H_2O_2$. Many studies have suggested that the ROS generation might be a markers for oxidative stress. As shown in Fig. 2A, H_2O_2 treatment for 1 hour was found to markedly increased intracellular ROS by maximum of $626.1 \pm 10.2\%$ as compared to that those of the controls ($p < 0.05$). Remarkably, ursolic acid exposure was found to significantly decrease the H_2O_2 -induced dichlorofluorescein fluorescence in dose-dependent manner ($p < 0.05$) (Fig. 2B).

Furthermore, ALP activity in H_2O_2 -treated MG-63 cells was found to be markedly decreased by 30.0% as compared to that of the control ($p < 0.01$). However, we

observed that the addition of $5 \mu M$ ursolic acid significantly increased the ALP activity at concentration for 10 days in H_2O_2 -treated osteoblasts ($p < 0.05$) (Fig. 2C).

3. Ursolic acid-mediated protective action involves immediate early response 3/nuclear factor erythroid 2-related factor 2 pathway

Accumulated evidences have showed that Nrf2 is a transcriptional factor that plays a key role in antioxidant response element-mediated induction of phase II detoxifying and antioxidant enzymes. To further investigate the molecular mechanism of ursolic acid against oxidative stress in osteoblasts, we measured the protein expression levels of IER3/Nrf2-related genes (IER3, NRF2, and HO-1) by immunoblotting. As shown in Fig. 3, the expression levels of Nrf2 and HO-1 were found to be increased in MG-63 cells when treated with ursolic acid as compared to their expression levels in the controls and H_2O_2 alone-treated control cells. Conversely, the expression levels of IER3 was observed lower than the expression level in H_2O_2 alone-treated control cells. In addition, ursolic acid was found to up-regulate HO-1 expression in a concentration-dependent manner. Thus, our results indicated that ursolic acid has a protective effect against H_2O_2 -induced oxidative stress by inhibiting IER3 expression, and up-regulating the expression of Nrf2 and HO-1 in osteoblasts.

Fig. 2. Inhibitory effects of ursolic acid on reactive oxygen species production and osteoblastic differentiation by H_2O_2 . (A) Cells were treated with $100 \mu M H_2O_2$, and dichlorofluorescein fluorescence intensity was confirmed for 60 minutes. (B) and (C) cells were pretreated with ursolic acid ($0.5 \sim 10 \mu M$) for 2 hours and then incubated with H_2O_2 . Values are mean $\pm$ standard deviation of three experiments. * $p < 0.05$ and ** $p < 0.01$ vs. control group, *** $p < 0.05$ vs. H_2O_2 alone). ALP: alkaline phosphatase.

Fig. 3. Involvement of immediate early response 3 (IER3)/nuclear factor erythroid 2-related factor 2 (Nrf2) pathway in the protective effects of ursolic acid on the H₂O₂-induced inhibition of osteoblastic differentiation of MG-63. Protein expression of signaling pathway-related regulators IER3, Nrf2, and heme oxygenase-1 (HO-1) were analyzed by western blot assay. The data presented are representative of three independent experiments.

Discussion

Oxidative stress in the periodontal microenvironment is a reactive, cellular defense mechanism that produces the energy required for physiological activity, and to destroys pathogens and their components¹⁸. Excessive endogenous production of ROS in periodontitis leads to an imbalance in the antioxidant defense system. Increase in ROS levels in lesions might indicate that they contribute in the impairment of periodontal tissue¹⁹. A recent study using dental stem cells demonstrated that oxidative stress conditions negatively influences cellular ability to repair and regenerate periodontal tissue²⁰.

Periodontitis is a chronic inflammatory disease that leads to bone impairment via pathogen complex and various stimulants and induces oxidative damage during chronic inflammation⁹. As imbalance in bone resorption and formation plays an important role in the disease progression, periodontal hard tissue regeneration studies have gained much attention. However, to our knowledge, there has been no definite recommended treatment for bone tissue regeneration in periodontal lesions yet.

Historically, the phytochemical and pharmacological activities of natural compounds from traditional medicines have been investigated in the early stages of drug discovery. In addition, ursolic acid has been shown to stimulate in vitro differentiation of osteoblastic cells, MC3T3-E1²¹ and also proven to have in vivo bone-forming activity in a mouse calvarial bone-formation model²². Presently, it has

been demonstrated that ursolic acid may play an important role in bone repair and regeneration.

The present study proposes that H₂O₂-induced production of ROS in MG-63 osteoblasts could be partially inhibited by ursolic acid. In this study, cells were treated with ursolic acid (0 μM, 0.5 μM, 1 μM, 5 μM, or 10 μM) in OS medium for 10 days to optimize the proliferation of MG-63 osteoblasts. Our results showed that ursolic acid promoted the proliferation and osteoblastic differentiation of MG-63 cells in a dose-dependent manner. Moreover, ursolic acid was found to significantly attenuate the inhibition of osteoblastic differentiation under a high oxidative stress condition. Our results are consistent with the previous findings on stimulating osteoblast differentiation and inducing new bone formation in bone²².

IER3, also known as immediate early gene X-1, expression is induced by a wide range of stimuli, such as growth factors, cytokines, ionizing radiation, and viral infections^{23,24}. IER3 suppresses the production of ROS and protects cells from apoptosis under stress conditions. Transcription factor Nrf2, has been shown to exhibit resistance to oxidative stress and promote of bone formation²⁵. Previous reports have revealed that Nrf2 regulates the expression of various proteins such as phase II detoxifying enzymes. Additionally, it has been reported that HO-1 is a downstream protein regulated by Nrf2, and associated with antioxidant genes that protect cells from various injuries via their anti-inflammatory effects and regulate disease progression²⁶. Furthermore, overexpression

of NRF2 in mesenchymal stem cells is reported to enhance osteoblastic differentiation and prevent apoptosis under oxidative stress²⁷⁾. Regulation of Nrf2 activation has been reported to enhance oxidative stress during chronic inflammation by IER3. This indicate that a novel mechanism regulates Nrf2 regulation by which IER3 exhibits its stress adaptability and tumor-suppressive activity²⁸⁾. Interestingly, the current study demonstrated that ursolic acid induces protein expression of the stress response genes, *HO-1* through multifunctional stress response gene; *IER3*, and transcription factor Nrf2.

In conclusion, the present study suggests that ursolic acid promotes osteoblastic differentiation via activation of defense mechanisms. These findings provide new evidence for the potential clinical applications of ursolic acid in oxidative stress conditions associated with periodontal disease.

Notes

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Ethical approval

This article is not necessary for IRB screening because it is an experimental paper using commercially available cells.

ORCID

Sang-im Lee, <https://orcid.org/0000-0003-2635-6876>

Acknowledgements

This research was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (No. 2018R1D1A1B07051253).

References

- Pizzino G, Irrera N, Cucinotta M, et al.: Oxidative stress: harms and benefits for human health. *Oxid Med Cell Longev* 2017: 8416763, 2017. <https://doi.org/10.1155/2017/8416763>
- Espinosa-Diez C, Miguel V, Mennerich D, et al.: Antioxidant responses and cellular adjustments to oxidative stress. *Redox Biol* 6: 183-197, 2015. <https://doi.org/10.1016/j.redox.2015.07.008>
- Mittal M, Siddiqui MR, Tran K, Reddy SP, Malik AB: Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal* 20, 1126-1167, 2014. <https://doi.org/10.1089/ars.2012.5149>
- Aruoma OI: Free radicals, oxidative stress, and antioxidants in human health and disease. *J Am Oil Chem Soc* 75: 199-212, 1998. <https://doi.org/10.1007/s11746-998-0032-9>
- Manolagas SC: From estrogen-centric to aging and oxidative stress: a revised perspective of the pathogenesis of osteoporosis. *Endocr Rev* 31: 266-300, 2010. <https://doi.org/10.1210/er.2009-0024>
- D'Aiuto F, Nibali L, Parkar M, Patel K, Suvar J, Donos N: Oxidative stress, systemic inflammation, and severe periodontitis. *J Dent Res* 89: 1241-1246, 2010. <https://doi.org/10.1177/0022034510375830>
- Bostanci N, Emingil G, Saygan B, et al.: Expression and regulation of the NALP3 inflammasome complex in periodontal diseases. *Clin Exp Immunol* 157: 415-422, 2009. <https://doi.org/10.1111/j.1365-2249.2009.03972.x>
- Nazir MA: Prevalence of periodontal disease, its association with systemic diseases and prevention. *Int J Health Sci (Qassim)* 11: 72-80, 2017.
- Oktay S, Chukkapalli SS, Rivera-Kweh MF, Velsko IM, Holliday LS, Kesavalu L: Periodontitis in rats induces systemic oxidative stress that is controlled by bone-targeted antiresorptives. *J Periodontol* 86: 137-145, 2015. <https://doi.org/10.1902/jop.2014.140302>
- Zheng DH, Han ZQ, Wang XX, Ma D, Zhang J: Erythropoietin attenuates high glucose-induced oxidative stress and inhibition of osteogenic differentiation in periodontal ligament stem cell (PDLSCs). *Chem Biol Interact* 305: 40-47, 2019. <https://doi.org/10.1016/j.cbi.2019.03.007>
- Wang Y, Wu H, Shen M, Ding S, Miao J, Chen N: Role of human amnion-derived mesenchymal stem cells in promoting osteogenic differentiation by influencing p38 MAPK signaling in lipopolysaccharide -induced human bone marrow mesenchymal stem cells. *Exp Cell Res* 350: 41-49, 2017. <https://doi.org/10.1016/j.yexcr.2016.11.003>
- Dahiya P, Kamal R, Gupta R, Bhardwaj R, Chaudhary K, Kaur S: Reactive oxygen species in periodontitis. *J Indian*

- Soc Periodontol 17: 411-416, 2013.
<https://doi.org/10.4103/0972-124X.118306>
13. Xing Y, Zhang Y, Jia L, Xu X: Lipopolysaccharide from *Escherichia coli* stimulates osteogenic differentiation of human periodontal ligament stem cells through Wnt/ β -catenin-induced TAZ elevation. *Mol Oral Microbiol* 34: 1-13, 2019. <https://doi.org/10.1111/omi.12249>
14. Peng M, Qiang L, Xu Y, Li C, Li T, Wang J: Inhibition of JNK and activation of the AMPK-Nrf2 axis by corosolic acid suppress osteolysis and oxidative stress. *Nitric Oxide* 82: 12-24, 2019. <https://doi.org/10.1016/j.niox.2018.11.002>
15. Kanzaki H, Shinohara F, Kajiya M, Kodama T: The Keap1/Nrf2 protein axis plays a role in osteoclast differentiation by regulating intracellular reactive oxygen species signaling. *J Biol Chem* 288: 23009-23020, 2013.
<https://doi.org/10.1074/jbc.M113.47854>
16. Liu J: Oleanolic acid and ursolic acid: research perspectives. *J Ethnopharmacol* 100: 92-94, 2005.
<https://doi.org/10.1016/j.jep.2005.05.024>
17. Sultana N: Clinically useful anticancer, antitumor, and antiwrinkle agent, ursolic acid and related derivatives as medicinally important natural product. *J Enzyme Inhib Med Chem* 26: 616-642, 2011.
<https://doi.org/10.3109/14756366.2010.546793>
18. Bullón P, Román-Malo L, Marín-Aguilar F, et al.: Lipophilic antioxidants prevent lipopolysaccharide-induced mitochondrial dysfunction through mitochondrial biogenesis improvement. *Pharmacol Res* 91: 1-8, 2015.
<https://doi.org/10.1016/j.phrs.2014.10.007>
19. Aboodi GM, Goldberg MB, Glogauer M: Refractory periodontitis population characterized by a hyperactive oral neutrophil phenotype. *J Periodontol* 82: 726-733, 2011.
<https://doi.org/10.1902/jop.2010.100508>
20. Kook SH, Lee D, Cho ES, et al.: Activation of canonical Wnt/ β -catenin signaling inhibits H₂O₂-induced decreases in proliferation and differentiation of human periodontal ligament fibroblasts. *Mol Cell Biochem* 411: 83-94, 2016.
<https://doi.org/10.1007/s11010-015-2570-4>
21. Li Q, Fan YS, Gao ZQ, Fan K, Liu ZJ: Effect of Fructus *Ligustri Lucidi* on osteoblastic like cell-line MC3T3-E1. *J Ethnopharmacol* 170: 88-95, 2015.
<https://doi.org/10.1016/j.jep.2015.05.013>
22. Lee SU, Park SJ, Kwak HB, Oh J, Min YK, Kim SH: Anabolic activity of ursolic acid in bone: stimulating osteoblast differentiation in vitro and inducing new bone formation in vivo. *Pharmacol Res* 58: 290-296, 2008.
<https://doi.org/10.1016/j.phrs.2008.08.008>
23. Arlt A, Minkenberg J, Kocs B, et al.: The expression of immediate early gene X-1 (IEX-1) is differentially induced by retinoic acids in NB4 and KG1 cells: possible implication in the distinct phenotype of retinoic acid-responsive and -resistant leukemic cells. *Leukemia* 18: 1646-1655, 2004.
<https://doi.org/10.1038/sj.leu.2403481>
24. Taddeo B, Esclatine A, Zhang W, Roizman B: The stress-inducible immediate-early responsive gene IEX-1 is activated in cells infected with herpes simplex virus 1, but several viral mechanisms, including 3' degradation of its RNA, preclude expression of the gene. *J Virol* 77: 6178-6187, 2003.
<https://doi.org/10.1128/jvi.77.11.6178-6187.2003>
25. Liu S, Fang T, Yang L, Chen Z, Mu S, Fu Q: Gastrodin protects MC3T3-E1 osteoblasts from dexamethasone-induced cellular dysfunction and promotes bone formation via induction of the NRF2 signaling pathway. *Int J Mol Med* 41: 2059-2069, 2018. <https://doi.org/10.3892/ijmm.2018.3414>
26. Chen XL, Dodd G, Thomas S, et al.: Activation of Nrf2/ARE pathway protects endothelial cells from oxidant injury and inhibits inflammatory gene expression. *Am J Physiol Heart Circ Physiol* 290: H1862-H1870, 2006.
<https://doi.org/10.1152/ajpheart.00651.2005>
27. Yuan Z, Zhang J, Huang Y, et al.: NRF2 overexpression in mesenchymal stem cells induces stem-cell marker expression and enhances osteoblastic differentiation. *Biochem Biophys Res Commun* 491: 228-235, 2017.
<https://doi.org/10.1016/j.bbrc.2017.07.083>
28. Stachel I, Geismann C, Aden K, et al.: Modulation of nuclear factor E2-related factor-2 (Nrf2) activation by the stress response gene immediate early response-3 (IER3) in colonic epithelial cells: a novel mechanism of cellular adaption to inflammatory stress. *J Biol Chem* 289: 1917-1929, 2014.
<https://doi.org/10.1074/jbc.M113.490920>